

УДК 616.64:616.9-002.828:582.282.23-085
DOI 10.24144/1998-6475.2022.55.23-30

BASIS FOR EMERGENCE OF SEBORRHEIC DERMATITIS

Vysochanska Vlasta, Koval Galina

*Uzhhorod National University. Medical Faculty. Microbiology, virology, epidemiology department
with the course of infectious disease.*

Abstract. *Introduction.* Seborrheic dermatitis (SD) is commonly associated with presence of *Malassezia* fungi. As a part of microbiome *M. restricta*, *M. furfur*, *M. globosa* protects us from pathogenic bacteria and UV-light and can be found at any skin zone of our body, except palms and feet. Hyperproduction of sebum, inflammation, and itching as symptoms of SD also are caused by many diseases related with endocrine, neuro-, immune, and psychological disorders. It is impossible to unambiguously identify the cause of inflammation of the skin in areas of large accumulation of sebaceous glands. Therefore, current treatment of patients with seborrheic dermatitis requires a personalized and predictive approach including skin phenotype, somatic and mental comorbid disease.

Objective: to determine the factors that create the prerequisites for the development of dermatitis and overgrowth of *Malassezia* spp.

Materials and methods: Articles from Wiley online library, Cochrane library, Medline were analyzed.

Conclusions: SD should be considered as an early manifestation or visible symptom of somatic or mental illnesses. *Malassezia* colonization in sebaceous areas is an outcome, not a cause of SD.

Key words: *Malassezia* spp., seborrheic dermatitis, sebaceous gland.

Передумови для виникнення себорейного дерматиту

Височанська В.В., Коваль Г.М.

Резюме. *Вступ.* Себорейний дерматит (СД) часто асоціюють із присутністю на шкірі грибів роду *Malassezia*. Будучи частиною мікробіому *M. restricta*, *M. furfur*, *M. globosa* захищають від впливу УФ випромінювання та патогенних бактерій і колонізують практично всі ділянки шкіри, крім долонь та стоп. Надмірна секреція шкірного сала, запалення та свербіж, які прийнято вважати основними симптомами СД, частіше за все спричинені нейро-, ендокринними, імунними та психічними розладами. Неможливо точно встановити причину запалення шкіри у ділянках з великою кількістю сальних залоз. Тому сучасне лікування пацієнтів із себорейним дерматитом повинно мати персоналізований і предиктивний підхід і враховувати фенотип шкіри, соматичні та психічні коморбідні розлади.

Метою дослідження було встановити фактори, які створюють передумови для виникнення себорейного дерматиту та надмірного росту *Malassezia* spp.

Матеріали та методи: аналіз оригінальних досліджень із бібліотек Wiley online library, Cochrane library, Medline.

Висновки. Себорейний дерматит варто розглядати як прояви маніфестації чи єдині видимі симптоми деяких соматичних та психічних захворювань. Колонізація запалених ділянок шкіри грибами *Malassezia* spp. є наслідком адаптації гриба до нових умов існування, а не причиною виникнення себорейного дерматиту.

Ключові слова: *Malassezia* spp., себорейний дерматит, сальна залоза.

Introduction

It is believed that seborrheic dermatitis is often caused by fungi of the genus *Malassezia*. The main approach in SD therapy are drugs with antifungal activity: azoles, hydroxypyridones, allylamines, selenium and pyrithione of zinc, carbonic anhydrases¹. But why corticosteroids¹, calcineurin inhibitors², cancer drugs³, statins⁴ help? In our opinion, *Malassezia* actively colonizes those areas of the skin where the

environment is most favorable. And the basis for emergences of seborrheic dermatitis is the excessive production of sebum and required pH. Therefore, for good control of symptoms, it is necessary to treat the cause, not to eliminate the fungus.

Main reason of skin inflammation, desquamation and pruritus is based on idea of pathological activity of saprophytic fungi of the genus *Malassezia* (formerly known as *Pityrosporum*)⁵ such as



*Malassezia restricta*⁶, *M. furfur*⁷, *M. globosa*⁸. It is believed that *Malassezia* spp.:

- enhance the inflammatory response by inducing the synthesis of IL-1 β , IL-1 α , IL-8, but inhibit the synthesis of cytokines by keratinocytes,⁹
- synthesize metabolites that, acting on aryl-hydrocarbon receptors (Ah-receptors), impede the maturation of dendritic cells,¹⁰
- cause apoptosis of keratinocytes,¹¹
- violate skin homeostasis,¹²
- are associated with Paget's disease¹³ and carcinogenesis.¹⁴
- is capable of biofilm formation¹⁵ and symbiosis with *Staphylococcus aureus*¹⁶ and *Bacillus* spp.¹⁷

Malassezia have genes encoding secreted hydrolases (e.g., lipases, phospholipases, aspartyl proteases, and acid sphingomyelinases), but do not possess genes for fatty acid synthase¹⁸. Therefore, they are limited by their inability to synthesize lipids and are forced to survive in an environment with many sebaceous glands. *Malassezia* need both saturated (palmitic acid (PA) and unsaturated acids (oleic acid (OA) to eat and live. PA and OA have fungicidal¹⁹ and fungistatic effect²⁰ on *M. furfur*, and on *M. sympodialis* at the same time. We might assume that skin fatty metabolism and *Malassezia* colonization is a self-regulated system. Plus, not eaten oleic acid causes declension of epidermal differentiation, barrier defects and disorders of skin metabolism²¹ and protein metabolites of *Malassezia* protect from UV light.²² So is there a real need to get rid of fungus?

Oily and dry skin has always been associated with SD. This may indicate increased activity of the sebaceous glands and impaired function of the epidermal barrier. Instead, the decrease in lipid synthesis of sebocytes protects the epidermis from morphological changes caused by *Malassezia* and reduces dandruff. However, the role of sebum in the pathogenesis of SD remains controversial, since hyperactivity does not always cause dermatitis, and patients with SD may not have oily skin²³. Some authors even suggest using "dermatitis of the sebaceous areas" than "seborrheic dermatitis" as a more precise term²⁴.

The aim of the study

The aim of our study was to determine the factors that create the prerequisites for the development of dermatitis and overgrowth of *Malassezia* spp.

Materials and methods

Articles from Wiley online library, Cochrane library, Medline were analyzed.

Results

1. Features of the sebaceous glands.

The holocrine type of sebaceous gland secretion (SG) is characterized by the fact that the glandular cell fills with fat and dies. The secret has been excreted on the surface of the epithelium near the hair follicles. Most SGs are focused on the face (especially the T-zone), the back, and the chest. Sebum acts as a delivery system for antimicrobial lipids, antioxidants, pheromone generation. It also serves as a substrate for bacterial and yeast growth²⁵.

2. There are numerous studies describing various forms of regulation of the sebaceous glands²⁶⁻²⁹ and their effect on *Malassezia* spp. Let us look at some of them.

2.1 Endocrine regulation.

The hypothalamic-pituitary-adrenal system or axis (HPA) involves several neuro- and steroid hormones. Corticotropin-releasing factor or corticoliberin stimulates the release of adrenocorticotrophic hormone (ACTH). HPA has shared receptors with the melanocortin system that regulate pigmentation and homeostasis. In addition to systemic regulation, the skin itself synthesizes some components of the HPA axis³⁰. Therefore, it can be assumed that the sebaceous glands are regulated not only by systemic but also by autocrine impact of the skin.

Corticotropin acts directly on sebocytes. The introduction of ACTH to rats leads to enlargement of the sebaceous glands and intensification of their secretion. Interestingly, this effect is evident even in rats with removed adrenals. Therefore, ACTH can stimulate the sebaceous glands regardless of central regulation. In addition, ACTH is independent of the presence testicles and has no synergistic effect with testosterone. Alpha melanocyte stimulating hormone (α -MSG) also independently stimulates sebum secretion but exhibits a synergistic effect when is used with testosterone³¹.

Prolactin is a pituitary neurohormone that regulates lactation and is dependent on the menstrual cycle. High levels of prolactin during pregnancy are reduced by childbirth but may be stimulated by breastfeeding. Interestingly, such hormone vacillations correlate with sebaceous gland secretion. It is believed that mom's aroma controls the behavior and reflexes of the baby³².

Increased secretion is explained by the presence of prolactin receptors on sebocytes³³.

Abnormally high levels of prolactin can be observed in Parkinson's disease (PD) as well as excessive sebum production. Parkinson is a disease of dopaminergic neurons with dopamine deficiency. Seborrhea and hyperprolactinemia can be treated by restoring normal levels of dopamine through the administration of L-DOPA or using dopaminergic agonists³⁴.

Phenylketonuria (PKU) is another hypodopaminergic condition characterized by hyperprolactinemia and seborrhea and might be treated with L-DOPA³⁵.

Glucocorticoids enhance hair growth, stimulate sebocyte maturation and cause skin atrophy by acting on fibroblasts. Growth factors that exacerbate sebaceous gland diseases have a synergistic effect³⁶.

Growth hormone (GR) is produced mainly by somatotrophic cells of the anterior pituitary gland. Acromegaly, as a manifestation of excessive GR in adulthood, is accompanied by increased sebum secretion. Conversely, GH deficiency is associated with abnormally low sebum secretion, which can be normalized after the use of substitution therapy. This conclusion was confirmed in a study demonstrating the presence of GR receptors on sebocytes³⁷.

Insulin-like growth factor has long been associated with weight loss diets. Low carbohydrate diets are known to have a positive effect in the treatment of acne. Hormone levels have been found to be positively correlated with sebum production³⁸.

Thyrotropic hormone (TTH), **triiodothyronine** (T3) and **thyroxine** (T4) are important hormones that regulate growth and development. It has been observed that hypothyroidism and thyroidectomy are associated with abnormally low sebum secretion. L-thyroxine replacement therapy partially normalizes the function of the glands, and thyrotoxicosis (hyperthyroidism) increases it excessively³⁹. It tends to think that thyroid hormones act independently of androgens. TTH and T4 enhance the activity sebaceous glands even in castrated animals, and additional stimulation with testosterone does not produce a synergistic effect. TTH stimulation of animals with thyroid resection does not affect the sebaceous glands, which confirms the need in thyroid hormones. SG hyperstimulation is possible due to the presence of specific T3 receptors on their surface⁴⁰.

Androgens such as dehydroepiandrosterone (DHEA), androstenedione (A-dione) and testosterone stimulate sebum secretion in humans and increase gland size. The action of the active forms of androgens is mediated by their binding to nuclear androgen receptors. The absence of functional androgen receptors, for example, in general androgen insensitivity syndrome, impedes the action of androgens on skin appendages⁴¹.

Therefore, skin synthesis and degradation of active androgens are noteworthy. It is known that precursors of active forms in tissues, such as DHEA DHEA and A-dione, are mainly synthesized by the glands, and testosterone and 5 α -dihydrotestosterone (5 α -DHT) in women are synthesized on the periphery. Androgens alone cannot affect sebocyte differentiation. They are activated by peroxisome proliferator-activated receptor (PPARs) ligands. Combining local hyperproduction of active androgens or an increase in blood 5 α -DHT levels at normal levels of DHEA, A-dione, and testosterone may be the cause of sebum hyperproduction⁴².

Estrogen β -receptors are present in keratocytes, melanocytes, fibroblasts and sebocytes, α -receptors have been found only in sebocytes in vitro. Estrogens reduce the activity of the sebaceous glands and are direct testosterone antagonists⁴³.

2.2. Neuropeptide regulation.

Nerve fibers due to release neuropeptides and neurotransmitters control epidermal proliferation, apoptosis, wound healing, inflammation, and melanogenesis. Eccrine glands and hair follicles are surrounded by a thick net of nerve endings. The presence of direct innervation of the sebaceous glands is noticeable in patients with partial paralysis of the face. The plegic side has increased sebum secretion and is more prone to acne⁴⁴. The opposite effect is achieved by using topical antimuscarinic agents and botulinum toxin-A (Botox), which is used to reduce acne and sebum in humans⁴⁵. Unfortunately, no study can accurately demonstrate the size and localization of nerve fibers that directly innervate the sebaceous glands. The possibility of indirect effects on the sebaceous glands by innervation of their ancestors near the hair follicles (HF) is considered. HF have a vast neural network that consists of a cluster of perifollicular and longitudinal nerve endings. The number of nerve fibers near the HF is demonstrated by the dependence of the allocation of fat on the hair growth cycle⁴⁶.



Acetylcholine directly affects the expression of cholinergic receptors in SG by stimulating muscarinic m2AChR and nicotinic receptors nAChR α 7, nAChR α 10 and nAChR β 4⁴⁷. The increased sebum production on the face is observed after the application of acetylcholine and is directly proportional to the dose. Stimulation by non-neural sources of acetylcholine in the skin also should not be denied⁴⁸.

Adrenaline, as the major neurotransmitter of the sympathetic nervous system, does not affect the level of sebum secretion, which has been demonstrated by β -receptors blockade with Propranolol. And the sebaceous glands have neither β 1- nor β 2-adrenergic receptors⁴⁹.

Substance P (CP) is a neuropeptide of the tachykinin genus that modulates nociception, inflammation, and vasodilatation. CP-sensitive connective tissue fibers were found around the sebaceous glands, but only in patients with acne. Mice treated with CP developed expanded sebaceous glands and increased fat production. In humans it has also caused an increase of the level of interleukins and tumor necrosis factor. The inflammatory reaction disappeared with the introduction of dexamethasone. However, proven involvement of CP in the pathogenesis of inflammation is only available in patients with acne⁵⁰.

Neutrophins are proteins that regulate axonal growth and trophic supply of neurons. The sebaceous glands have receptors for neurotrophin -3 and nerve growth factor (NGF). Stress and inflammatory skin diseases induce NGF synthesis as opposed to healthy skin patients. Nevertheless, the role of neurotrophins in the regulation of skin fat production remains unknown but suggests the existence of a paracrine or autocrine mechanism of neurotrophin regulation of sebocytes⁵¹. An alternative version is an indirect action of neurotrophins by stimulating the growth of nerve fibers around the sebaceous cells of the progenitors of SG. In the experiment on skin, denervation led only to a slight decrease of hair growth, which may be an evidence of autocrine regulation rather than a direct enhancement of innervation⁵².

Endocannabinoids are unsaturated fatty acids. They are synthesized by the epithelium and modulate nociceptive and temperature sensitivity of skin nerves. Sebocytes have cannabinoid receptors and TRP channels (transient receptor potential channels). Activated endocannabinoid receptors modulate thermoregulation and

enhance pain and itching sensations. Endogenous stimulation of these receptors leads to increased lipogenesis⁵³. The effect of increasing endocannabinoid levels and phytocannabinoid (cannabigerol, cannabigerovarin) has been proposed as a treatment for inflammatory conditions accompanied by dry skin. In contrast to phytocannabinoids, cannabidiol, cannabichromen and cannabidivarin combine lipostatic, antiproliferative and anti-inflammatory effects and have the potential to treat acne⁵⁴.

Calcitonin gene-bound peptide (CGRP) is a calcitonin receptor that is also present on sebocytes⁵⁵.

2.3. Trait of the host.

Immunity, according to the literature, is a key weapon in the fight against *Malassezia* infections. Immunocompromised patients have seborrheic dermatitis much more often than healthy ones⁵⁶. Skin affected by dermatitis is accompanied by increased immunoreactivity and high levels of IL-1 α , IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12, β -defensin, TNF α and IFN γ . Also, due to the increased degranulation of mast cells, there is a hyperproduction of histamine that accumulates in the epidermal layer. The DNA microarray test of patients with dandruff confirmed the activity of inflammatory genes and showed decreased activity of genes controlling lipid metabolism⁵⁷. Also, mice with activated 2C T-cell receptor had severe CD4 + and CD8 + T cell lymphopenia, and subsequently spontaneously developed a seborrheic skin phenotype. Similar abnormalities have been observed in HIV-positive patients, where it is known that the prevalence of SD is extremely high⁵⁶.

The microbiome of healthy skin differs from patients with seborrheic dermatitis. SD patients are mostly colonized by *Acinetobacter spp.*, *Staphylococcus spp.* and *Streptococcus sp.*⁵⁸. Among patients with dandruff, the most common are *Basidiomycota* (*Filobasidium spp.*) while among the healthy individuals *Ascomycota* (*Acremonium spp.*)⁵⁹. Innate lymphoid cells (ILCs) affect commensal bacteria by modulating sebaceous gland activity in mice. Pools of ILC are found in all parts of the skin to inhibit sebocyte proliferation. In their absence, the sebaceous glands enlarge and more actively produce PA and OA. PA acid has specific anti-Gram-positive agents' activity. It is possible that with an excessive amount of substrate commensal *Malassezia* overgrows and triggers an inflammatory response but depresses a bacterial growth.

Skin barrier defects are associated with exacerbations of SD. Dry skin in winter and maceration most often violate skin protection. *Malassezia* modulates skin function directly through mediators or indirectly activate the immune response. It has been described that eradication of some *Malassezia* spp. makes ceramides, sphingolipids, histamine, and inflammatory markers in the skin return to norma⁶⁰.

Conclusions

The survival of *Malassezia* depends on the rate of adaptation to sweat, sebum, acidity, immune response, humidity, and environmental

insolation, as well as the activity of other skin commensals - viruses, bacteria, microscopic fungi. *Malassezia* may trigger immune response but also protects our skin from UV-light, action of sebum and sweat and pathogenic bacteria. It is impossible to unambiguously identify the cause of inflammation of the skin in areas of large accumulation of sebaceous glands. Therefore, current treatment of patients with seborrheic dermatitis requires a personalized and predictive approach. Seborrheic dermatitis can be a marker of somatic diseases (see Table 1). *Malassezia* colonization in sebaceous areas is a consequence, not a cause of SD.

Table 1

Pathological conditions and disease having impact on seborrheic dermatitis

Diseases/ conditions	Sebum production	References	Article
Addison's disease	Sebum ↓	Goolamali SK (1974)	Sebum excretion and melanocyte-stimulating hormone in hypoadrenalism.
		Z Haut Geschlechtskr (1962)	Small-nodular sebaceous gland hypertrophy after cortisone treatment of Addison's disease.
Acromegaly	Sebum ↑	Resende M (2012)	Prevalence of dermatologic disorders in 15 patients with acromegaly.
		Borlu M (2012)	Acromegaly is associated with decreased skin transepidermal water loss and temperature, and increased skin pH and sebum secretion partially reversible after treatment.
Atopic dermatitis	Sebum normal or ↓ <i>Malassezia</i> spp. +	Agrawal K (2018)	Effects of atopic dermatitis and gender on sebum lipid mediator and fatty acid profiles.
		Ramos-E-Silva M (2018)	Red face revisited: Endogenous dermatitis in the form of atopic dermatitis and seborrheic dermatitis.
Bardet-Biedl syndrome	Sebum ↑	Haws RM (2019)	Cutaneous findings in Bardet-Biedl syndrome.
Hypothyroidism	Sebum ↓	Zouboulis CC (1998)	The human sebocyte culture model provides new insights into development and management of seborrhoea and acne.
Hashimoto autoimmune thyroiditis	Sebum ↓	Goucha S (2002)	Association of seborrheic pemphigus and Hashimoto autoimmune thyroiditis. Apropos of a further case
Hyperhidrosis	Sebum ↑	Mozaffari AA (2009)	Localized seborrheic dermatitis with hyperhidrosis due to mite infestation in an Iranian cross-breed ram.
Depression	Sebum ?	Maietta G (1990)	Patients with mood depression have a high prevalence of seborrheic dermatitis.
Tardive dyskinesia	Sebum ↑	Reuven Sandyk (1990)	Role of androgens in tardive dyskinesia.
Schizophrenia	Sebum ↑	Vassileva S (1990)	Familial occurrence of multiple seborrheic keratoses and schizophrenia.
		George A (2018)	A perspective study of cutaneous manifestations in primary psychiatric disorders in a tertiary care hospital.



Continuation of Table

Psoriasis	Sebum ↓ Malassezia spp. norma	Takahashi H (2014)	Defective barrier function accompanied by structural changes of psoriatic stratum corneum.
		Takemoto A (2015)	Molecular characterization of the skin fungal microbiome in patients with psoriasis.
Paget's disease	Sebum ↓ Malassezia spp. +	Shamsadini S (2006)	Surrounding ipsilateral eruptive seborrheic keratosis as a warning sign of intraductal breast carcinoma and Paget's disease.
		Sato Y (2019)	Malassezia-derived aryl hydrocarbon receptor ligands enhance the CCL20/Th17/soluble CD163 pathogenic axis in extra-mammary Paget's disease.
Parkinson's Disease	Sebum ↑ Malassezia spp. +	Trivedi DK (2019)	Discovery of Volatile Biomarkers of Parkinson's Disease from Sebum.
		Arsic Arsenijevic VS (2014)	A laboratory-based study on patients with Parkinson's disease and seborrheic dermatitis: the presence and density of Malassezia yeasts, their different species and enzymes production.
Phenylketonuria	Sebum ↑	Burton JL (1975)	An abnormality in sebaceous function in phenylketonuria.
Pregnancy	Sebum ↑	Burton JL (1970)	Effect of pregnancy on sebum excretion.
Dyslipidemia, obesity	No impact	Betul Imamoglu (2016)	Metabolic syndrome may be an important comorbidity in patients with seborrheic dermatitis.

REFERENCES

- Cohen SR, Gordon SC, Lam AH, Rosmarin D. Recalcitrant Seborrheic Dermatitis Successfully Treated With Apremilast. J Cutan Med Surg. 2020 Jan/Feb;24(1):90-91.
- Guenther L, Lynde C, Poulin Y. Off-Label Use of Topical Calcineurin Inhibitors in Dermatologic Disorders. J Cutan Med Surg. 2019 Sep/Oct;23(4_suppl):27S-34S.
- Retrosi C, Vezzoni R, Conforti C, Corneli P, Magaton-Rizzi G, Zalaudek I, Di Meo N. Targetoid erythema surrounding multiple seborrheic keratoses induced by chemotherapy with gemcitabine. Acta Dermatovenol Alp Pannonica Adriat. 2019 Dec;28(4):185-186.
- Sobhan M, Gholampoor G, Firozian F, Mohammadi Y, Mehrpooya M. Comparison of efficacy and safety of atorvastatin 5% lotion and betamethasone 0.1% lotion in the treatment of scalp seborrheic dermatitis. Clin Cosmet Investig Dermatol. 2019 Apr 29;12:267-275.
- Borda LJ, Wikramanayake TC. Seborrheic Dermatitis and Dandruff: A Comprehensive Review. J Clin Investig Dermatol. 2015 Dec;3(2):10.13188/2373-1044.1000019.
- Del Prete S, Vullo D, Ghobril C, Hitce J, Clavaud C, Marat X, et al. Cloning, Purification, and Characterization of a β -Carbonic Anhydrase from *Malassezia restricta*, an Opportunistic Pathogen Involved in Dandruff and Seborrheic Dermatitis. Int J Mol Sci. 2019 May 17;20(10):2447.
- Vest BE, Krauland K. Malassezia Furfur. 2021 Nov 7. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. PMID:31971731.
- Prohic A, Jovovic Sadikovic T, Krupalija-Fazlic M, Kuskunovic-Vlahovljak S. Malassezia species in healthy skin and in dermatological conditions. Int J Dermatol. 2016 May;55(5):494-504.
- Kistowska M, Fenini G, Jankovic D, et al. Malassezia yeasts activate the NLRP3 inflammasome in antigen-presenting cells via Syk-kinase signaling. Exp Dermatol. 2014 Dec; 23(12):884-9.
- Buommino E, Baroni A, Papulino C, et al. Malassezia pachydermatis up-regulates AhR related CYP1A1 gene and epidermal barrier markers in human keratinocytes. Med Mycol. 2018; 56(8):987-993.
- Krämer HJ, Podobinska M, Bartsch A, et al. Malassezin, a novel agonist of the aryl hydrocarbon receptor from the yeast Malassezia furfur, induces apoptosis in primary human melanocytes. Chem-biochem. 2005;6(5):860-5.
- Mexia N, Koutrakis S, He G, et al. A Biomimetic, One-Step Transformation of Simple Indolic Compounds to Malassezia-Related Alkaloids with High AhR Potency and Efficacy. Chem Res Toxicol. 2019;32(11):2238-2249.
- Sato Y, Fujimura T, Tanita K, et al. Malassezia-derived aryl hydrocarbon receptor ligands enhance the CCL20/Th17/soluble CD163 pathogenic axis in extra-mammary Paget's disease. Exp Dermatol. 2019;28(8):933-9.

14. Gaitanis G, Velegraki A, Magiatis P, et al. Could *Malassezia* yeasts be implicated in skin carcinogenesis through the production of aryl-hydrocarbon receptor ligands? *Med Hypotheses*. 2011;77(1):47-51.
15. Ianiri G, Heitman J, Scheynius A. The Skin Commensal Yeast *Malassezia globosa* Thwarts Bacterial Biofilms to Benefit the Host. *J Invest Dermatol*. 2018;138(5):1026-9.
16. Tamer F, Yuksel ME, Sarifakioglu E, et al. *Staphylococcus aureus* is the most common bacterial agent of the skin flora of patients with seborrheic dermatitis. *Dermatol Pract Concept*. 2018;8(2):80-4.
17. Sally G, Grimshaw, Adrian M. Smith, David S. Arnold, et al. The diversity and abundance of fungi and bacteria on the healthy and dandruff affected human scalp. *PLoS One*. 2019;14(12): e0225796.
18. Xu J, Saunders CW, Hu P, et al. Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proc Natl Acad Sci USA*. 2007;104(47):18730-5.
19. Triana S, de Cock H, Ohm RA, Danies G, Wösten HAB, Restrepo S, González Barrios AF, Celis A. Lipid metabolic versatility in *Malassezia* spp. yeasts studied through metabolic modeling. *Front. Microbiol*. 2017;8:1-18.
20. Kaneko T, Makimura K, Onozaki M, Ueda K, Yamada Y, Nishiyama Y, Yamaguchi H. Vital growth factors of *Malassezia* species on modified CHROMagar Candida. *Med. Mycol*. 2005; 43:699-704.
21. Wu G, Zhao Z, Li C, et al. Genus-Wide Comparative Genomics of *Malassezia* Delineates Its Phylogeny, Physiology, and Niche Adaptation on Human Skin. *PLoS Genet*. 2015;11(11): e1005614.
22. Mayser P, Stapelkamp H, Krämer H, et al. Pityriolactone- a new fluorochrome from the tryptophan metabolism of *M. alassezia furfur*. *Antonie Van Leeuwenhoek*. 2003;84:185-191.
23. Loing E, Lamarque E, Borel M. New targets in the battle against dandruff. *J Cosmet Sci*. 2017 Jan/Feb;68(1):107-113.
24. Shi VY, Leo M, Hassoun L, et al. Role of sebaceous glands in inflammatory dermatoses. *J Am Acad Dermatol*. 2015;73(5):856-63.
25. De Luca C, Valacchi G. Surface lipids as multifunctional mediators of skin responses to environmental stimuli. *Mediators Inflamm*. 2010;2010:321494.
26. Wikramanayake TC, Borda LJ, Miteva Met, et al. Seborrheic dermatitis-Looking beyond *Malassezia*. *Exp Dermatol*. 2019;28(9):991-1001.
27. Clayton RW, Langan EA, Ansell DM, et al. Neuroendocrinology and neurobiology of sebaceous glands. *Biol Rev Camb Philos Soc*. 2020.
28. Kamil Leis, Ewelina Mazur, et al. Endocrine systems of the skin. *Postepy Dermatol Alergol*. 2019;36(5):519-523.
29. Shi VY, Leo M, Hassoun L, et al. Role of sebaceous glands in inflammatory dermatoses. *J Am Acad Dermatol*. 2015;73(5):856-63.
30. Slominski AT, Zmijewski MA, Skobowiat C, Zbytek B, Slominski RM, Steketee JD. Sensing the environment: regulation of local and global homeostasis by the skin's neuroendocrine system. *Adv Anat Embryol Cell Biol*. 2012;212:v, vii, 1-115.
31. Thody AJ, Cooper MF, Bowden PE, Meddis D, Shuster S. Effect of alpha-melanocyte-stimulating hormone and testosterone on cutaneous and modified sebaceous glands in the rat. *J Endocrinol*. 1976 Dec;71(3):279-88.
32. Henderson CA, Taylor J, Cunliffe WJ. Sebum excretion rates in mothers and neonates. *Br J Dermatol*. 2000 Jan;142(1):110-1.
33. Langan EA, Hinde E, Paus R. Prolactin as a candidate sebotrop(h)ic hormone? *Exp Dermatol*. 2018 Jul;27(7):729-736.
34. Burton JL, Cartledge M, Shuster S. Effect of L-dopa on the seborrhoea of Parkinsonism. *Br J Dermatol*. 1973 May;88(5):475-9.
35. Juhász E, Kiss E, Simonova E, Patócs A, Reismann P. Serum prolactin as a biomarker for the study of intracerebral dopamine effect in adult patients with phenylketonuria: a cross-sectional monocentric study. *Eur J Med Res*. 2016 May 11;21(1):22.
36. Slominski A, Wortman J, Luger T, et al. Corticotropin releasing hormone and proopiomelanocortin involvement in the cutaneous response to stress. *Physiol Rev*. 2000;80:979-1020.
37. Borlu M, Karaca Z, Yildiz H, Tanriverdi F, Demirel B, Elbuken G, et al. Acromegaly is associated with decreased skin transepidermal water loss and temperature, and increased skin pH and sebum secretion partially reversible after treatment. *Growth Horm IGF Res*. 2012 Apr;22(2):82-6.
38. Ji J, Zhang RH, Li HM, Guo Q, Zhang LL, Zhu J, Chen L. Correlations of SOX9 expression with serum IGF1 and inflammatory cytokines IL-1 α and IL-6 in skin lesions of patients with acne. *Eur Rev Med Pharmacol Sci*. 2018 May;22(9):2549-55.



39. Ahsan MK, Urano Y, Kato S, Oura H, Arase S. Immunohistochemical localization of thyroid hormone nuclear receptors in human hair follicles and in vitro effect of L-triiodothyronine on cultured cells of hair follicles and skin. *J Med Invest.* 1998 Feb;44(3-4):179-84.
40. Bodó E, Kromminga A, Bíró T, Borbíró I, Gáspár E, Zmijewski MA, et al. Human female hair follicles are a direct, nonclassical target for thyroid-stimulating hormone. *J Invest Dermatol.* 2009 May;129(5):1126-39.
41. İslamoğlu ZGK. Second-to-fourth digit ratio and seborrheic dermatitis in males: a cross-sectional study. *An Bras Dermatol.* 2019 Jul 29;94(3):327-30.
42. Imperato-Mcginley, J. Zhu Yuan-Shan. Androgen and male physiology the syndrome of 5 α -reductase-2 deficiency. *Molecular and Cellular Endocrinology.* 2002;198:50-59.
43. Thornton MJ, Taylor AH, Mulligan K, Al-Azzawi F, Lyon CC, O'Driscoll J, Messenger AG. Oestrogen receptor beta is the predominant oestrogen receptor in human scalp skin. *Exp Dermatol.* 2003 Apr;12(2):181-90.
44. Sudy E, Urbina F. Unilateral acne after facial palsy. *An Bras Dermatol.* 2018 Jun;93(3):441-442.
45. Min P, Xi W, Grassetti L, Trisliana Perdanasari A, Torresetti M, Feng S, et al. Sebum Production Alteration after Botulinum Toxin Type A Injections for the Treatment of Forehead Rhytides: A Prospective Randomized Double-Blind Dose-Comparative Clinical Investigation. *Aesthet Surg J.* 2015 Jul;35(5):600-10.
46. Füllgrabe A, Joost S, Are A, Jacob T, Sivan U, Haegbarth A, et al. Dynamics of Lgr6⁺ Progenitor Cells in the Hair Follicle, Sebaceous Gland, and Interfollicular Epidermis. *Stem Cell Reports.* 2015 Nov 10;5(5):843-855.
47. Kurzen H, Wessler I, Kirkpatrick CJ, Kawashima K, Grando SA. The non-neuronal cholinergic system of human skin. *Horm Metab Res.* 2007 Feb;39(2):125-35.
48. Kurzen H, Wessler I, Kirkpatrick CJ, Kawashima K, Grando SA. The non-neuronal cholinergic system of human skin. *Horm Metab Res.* 2007 Feb;39(2):125-35.
49. Steinkraus V, Mak JC, Pichlmeier U, Mensing H, Ring J, Barnes PJ. Autoradiographic mapping of beta-adrenoceptors in human skin. *Arch Dermatol Res.* 1996 Aug;288(9):549-53.
50. Lee WJ, Jung HD, Lee HJ, Kim BS, Lee SJ, Kim DW. Influence of substance-P on cultured sebocytes. *Arch Dermatol Res.* 2008 Jul;300(6):311-6.
51. Toyoda M, Nakamura M, Makino T, Kagoura M, Morohashi M. Sebaceous glands in acne patients express high levels of neutral endopeptidase. *Exp Dermatol.* 2002 Jun;11(3):241-7.
52. Adly MA, Assaf HA, Hussein MR. Expression pattern of p75 neurotrophin receptor protein in human scalp skin and hair follicles: Hair cycle-dependent expression. *J Am Acad Dermatol.* 2009 Jan;60(1):99-109.
53. Dobrosi N, Tóth BI, Nagy G, Dózsá A, Géczy T, Nagy L, et al. Endocannabinoids enhance lipid synthesis and apoptosis of human sebocytes via cannabinoid receptor-2-mediated signaling. *FASEB J.* 2008 Oct;22(10):3685-95.
54. Oláh A, Tóth BI, Borbíró I, Sugawara K, Szöllösi AG, Czifra G, et al. Cannabidiol exerts sebostatic and antiinflammatory effects on human sebocytes. *J Clin Invest.* 2014 Sep;124(9):3713-24.
55. Böhm M, Schiller M, Ständer S, Seltmann H, Li Z, Brzoska T, et al. Evidence for expression of melanocortin-1 receptor in human sebocytes in vitro and in situ. *J Invest Dermatol.* 2002 Mar;118(3):533-9.
56. Moreno-Coutiño G, Sánchez-Cárdenas CD, Bello-Hernández Y, et al. Isolation of *Malassezia* spp. in HIV-positive patients with and without seborrheic dermatitis. *An Bras Dermatol.* 2019;94(5):527-531.
57. Trznadel-Grodzka E, Błaszowski M, Rotsztejn H. Investigations of seborrheic dermatitis. Part I. The role of selected cytokines in the pathogenesis of seborrheic dermatitis. *Postepy Hig Med Dosw (Online).* 2012;66:843-847.
58. Tanaka A, Cho O, Saito C, et al. Comprehensive pyrosequencing analysis of the bacterial microbiota of the skin of patients with seborrheic dermatitis. *Microbiol Immunol.* 2016; 60(8):521-6.
59. Park HK, Ha MH, Park SG, et al. Characterization of the fungal microbiota (mycobiome) in healthy and dandruff-afflicted human scalps. *PLoS One.* 2012;7(2):e32847.
60. Grice EA, Dawson TL Jr. Host-microbe interactions: *Malassezia* and human skin. *Curr Opin Microbiol.* 2017;40:81-87.